



HAL
open science

Generalized Bernoulli numbers and a formula of Lucas

Victor H. Moll, Christophe Vignat

► **To cite this version:**

Victor H. Moll, Christophe Vignat. Generalized Bernoulli numbers and a formula of Lucas. The Fibonacci Quarterly, 2015, 53 (4), pp.349. hal-01322848

HAL Id: hal-01322848

<https://centralesupelec.hal.science/hal-01322848>

Submitted on 10 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GENERALIZED BERNOULLI NUMBERS AND A FORMULA OF LUCAS

VICTOR H. MOLL AND CHRISTOPHE VIGNAT

ABSTRACT. An overlooked formula of E. Lucas for the generalized Bernoulli numbers is proved using generating functions. This is then used to provide a new proof and a new form of a sum involving classical Bernoulli numbers studied by K. Dilcher. The value of this sum is then given in terms of the Meixner-Pollaczek polynomials.

1. INTRODUCTION

The goal of this paper is to provide a unified approach to two topics that have appeared in the literature. The first one is an expression for the generalized Bernoulli numbers $B_n^{(p)}$ defined by the exponential generating function

$$(1.1) \quad \sum_{n=0}^{\infty} B_n^{(p)} \frac{z^n}{n!} = \left(\frac{z}{e^z - 1} \right)^p.$$

For $n \in \mathbb{N}$, the coefficients $B_n^{(p)}$ are polynomials in p named after Nörlund in [1]. The first few are

$$(1.2) \quad B_0^{(p)} = 1, B_1^{(p)} = -\frac{1}{2}p, B_2^{(p)} = -\frac{1}{12}p + \frac{1}{4}p^2, B_3^{(p)} = \frac{1}{8}p^2(1 - p).$$

In his 1878 paper E. Lucas [4] gave the formula

$$(1.3) \quad B_n^{(p)} = \frac{(-1)^{p-1} n!}{(p-1)! (n-p)!} \beta^{n-p+1} (1 + \beta) \cdots (p-1 + \beta)$$

for $n \geq p$. This is a symbolic formula: to obtain the value of $B_n^{(p)}$, expand the expression (1.3) and replace β^j by the ratio B_j/j . Here B_j is the classical Bernoulli number $B_n = B_n^{(1)}$ in the notation from (1.1).

The second topic is an expression established by K. Dilcher [2] for the sums of products of Bernoulli numbers

$$(1.4) \quad S_N(n) := \sum \binom{2n}{2j_1, 2j_2, \dots, 2j_N} B_{2j_1} B_{2j_2} \cdots B_{2j_N},$$

Date: February 14, 2014.

1991 Mathematics Subject Classification. Primary 11B68, Secondary 33C45.

Key words and phrases. Generalized Bernoulli numbers, Meixner-Pollaczek polynomials, Dilcher identities, Nörlund polynomials.

where the sum is taken over all nonnegative integers j_1, \dots, j_N such that $j_1 + \dots + j_N = n$, and where

$$(1.5) \quad \binom{2n}{2j_1, 2j_2, \dots, 2j_N} = \frac{(2n)!}{(2j_1)! \cdots (2j_N)!}$$

is the multinomial coefficient and B_{2k} is the classical Bernoulli number. One of the main results of [2] is the evaluation

$$(1.6) \quad S_N(n) = \frac{(2n)!}{(2n-N)!} \sum_{k=0}^{\lfloor (N-1)/2 \rfloor} b_k^{(N)} \frac{B_{2n-2k}}{2n-2k},$$

where the coefficients $b_k^{(N)}$ are defined by the recurrence

$$(1.7) \quad b_k^{(N+1)} = -\frac{1}{N} b_k^{(N)} + \frac{1}{4} b_{k-1}^{(N-1)},$$

with $b_0^{(1)} = 1$ and $b_k^{(N)} = 0$ for $k < 0$ and for $k > \lfloor (N-1)/2 \rfloor$.

Lucas's original proof is recalled in Section 2. This section also contains an extension of Lucas's formula for $B_n^{(p)}$ to $0 \leq n \leq p-1$ in terms of the Stirling numbers of the first kind. A unified proof of the two formulas for $B_n^{(p)}$ based on generating functions is given in Section 3. Another proof of Lucas's formula, based on recurrences, is given in Section 4 and Section 5 contains a proof of

$$(1.8) \quad S_N(n) = \sum_{k=0}^N \frac{(2n)!}{(2n-k)!} 2^{-k} \binom{N}{k} B_{2n-k}^{(N-k)}$$

that expresses Dilcher's sum (1.4) explicitly in terms of the generalized Bernoulli numbers. Expressing this result in hypergeometric form leads to a formula for $S_N(n)$ in terms of the Meixner-Pollaczek polynomials

$$(1.9) \quad P_n^{(\lambda)}(x; \phi) = \frac{(2\lambda)_n}{n!} e^{2n\phi} {}_2F_1 \left(\begin{matrix} -n & \lambda + ix \\ 2\lambda \end{matrix} \middle| 1 - e^{-2i\phi} \right).$$

It is then established that the recurrence (1.7), provided by Dilcher in [2], is equivalent to the classical three-term relation for this orthogonal family of polynomials.

2. LUCAS'S THEOREM

In his paper [4], E. Lucas gave an expression for the generalized Bernoulli numbers $B_n^{(p)}$, for $n \geq p$. This section presents an outline of his proof and an extension of this expression for $B_n^{(p)}$ to the case $0 \leq n \leq p-1$. A proof based on generating functions is given in the next section. Lucas's formula uses the translation

$$(2.1) \quad \beta^n = \frac{B_n}{n}$$

coming from umbral calculus. Observe, for example, that

$$\begin{aligned} B_3^{(2)} &= \frac{(-1)^1 3!}{1! 1!} \beta^2 (1 + \beta) = -6(\beta^2 + \beta^3) \\ &= -6 \left(\frac{B_2}{2} + \frac{B_3}{3} \right) = -3B_2 = -\frac{1}{2} \end{aligned}$$

Observe also that the symbolic substitution (2.1) should be performed only *after* all the terms have been expanded. For example,

$$(2.2) \quad \beta^2(1 + \beta) = \beta^2 + \beta^3 = \frac{B_2}{2} + \frac{B_3}{3} = -\frac{1}{4}$$

but

$$(2.3) \quad \beta^2(1 + \beta) \neq \frac{B_2}{2} \left(1 + \frac{B_1}{1} \right) = \frac{1}{24}.$$

Theorem 2.1 (Lucas). *For $n \geq p$, the generalized Bernoulli numbers $B_n^{(p)}$ are given by*

$$(2.4) \quad B_n^{(p)} = \frac{(-1)^{p-1} n!}{(p-1)! (n-p)!} \beta^{n-p+1} (1 + \beta)(2 + \beta) \cdots (p-1 + \beta)$$

where, in symbolic notation,

$$(2.5) \quad \beta^n = \frac{B_n}{n}.$$

Proof. Lucas's argument begins with the identity

$$(2.6) \quad pB_n^{(p+1)} = (p-n)B_n^{(p)} - pnB_{n-1}^{(p)}$$

which follows directly from the identity for generating functions

$$(2.7) \quad x \frac{d}{dx} \left(\frac{x}{e^x - 1} \right)^p = p(1-x) \left(\frac{x}{e^x - 1} \right)^p - p \left(\frac{x}{e^x - 1} \right)^{p+1}.$$

Shifting n to $n-1$ it follows that

$$(2.8) \quad pB_{n-1}^{(p+1)} = (p-n+1)B_{n-1}^{(p)} - p(n-1)B_{n-2}^{(p)}.$$

Now multiplying (2.6) by $n(p+1)$ and (2.8) by $(p-n+1)$ leads to

$$\begin{aligned} p(p+1)B_n^{(p+2)} &= (p-n+1)(p-n)B_n^{(p)} - (p-n+1)(p+p+1)nB_{n-1}^{(p)} \\ &\quad + p(p+1)n(n-1)B_{n-2}^{(p)} \end{aligned}$$

and then, by the same methods, he produces

$$\begin{aligned} (p+2)(p+1)pB_n^{(p+3)} &= (p-n+2)(p-n+1)(p-n)B_n^{(p)} \\ &\quad - (p-n_2)(p-n+1)(p+p+1+p+2)nB_{n-1}^{(p)} \\ &\quad + (p-n+2)(p(p+1) + p(p+2) + (p+1)(p+2))n(n-1)B_{n-2}^{(p)} \\ &\quad - p(p+1)(p+2)n(n-1)(n-2)B_{n-3}^{(p)} \end{aligned}$$

and then, stating '*and so on*', concludes the proof. □

The following alternate proof of Lucas's theorem using generating functions requires an expression for $B_n^{(p)}$ in the range $0 \leq n \leq p-1$, of the kind given in (2.4). This cannot be obtained directly from (2.4). The Stirling numbers of the first kind $s_k^{(p)}$ are used to produce an equivalent formulation of $B_n^{(p)}$. These numbers are defined by the generating function

$$(2.9) \quad z(z-1)(z-2)\cdots(z-(p-1)) = \sum_{k=1}^p s_k^{(p)} z^k.$$

Then (2.4) may be written as

$$\begin{aligned} B_n^{(p)} &= \frac{(-1)^{p-1}}{(p-1)!} n(n-1)\cdots(n-(p-1)) \beta^{n-p} (-1)^p \sum_{k=1}^p s_k^{(p)} (-\beta)^k \\ &= -\frac{1}{(p-1)!} n(n-1)\cdots(n-(p-1)) \sum_{k=1}^p s_k^{(p)} (-1)^k \frac{B_{n-p+k}}{n-p+k}. \end{aligned}$$

Observe that the index n varies in the range $0 \leq n \leq p-1$, therefore the prefactor $n(n-1)\cdots(n-(p-1))$ always vanishes. On the other hand, all the summands are finite, except when $k = p-n$. Performing the translation from $(-\beta)^k$ to B_k/k for this specific index gives

$$-\frac{1}{(p-1)!} n(n-1)\cdots 1 \times (-1)(-2)\cdots(-(p-1-n)) s_{p-n}^{(p)} (-1)^{p-n} = \frac{s_{p-n}^{(p)}}{\binom{p-1}{n}}.$$

This gives:

Theorem 2.2. *The generalized Bernoulli numbers $B_n^{(p)}$, with $0 \leq n \leq p-1$ are given by*

$$(2.10) \quad B_n^{(p)} = \frac{s_{p-n}^{(p)}}{\binom{p-1}{n}}.$$

In fact, this is a classical result. It is, for example, a direct consequence of the identity

$$(2.11) \quad (z-1)(z-2)\cdots(z-p) = \sum_{\ell=0}^p \binom{p}{\ell} z^\ell B_{p-\ell}^{(p+1)}$$

which appears (unnumbered) in [5, p.149].

3. THE PROOF VIA GENERATING FUNCTION

The expressions for the generalized Bernoulli numbers given in (2.4) and (2.10) are now used to compute the generating function

$$(3.1) \quad G(z) = \sum_{n=0}^{\infty} B_n^{(p)} \frac{z^n}{n!}$$

and to show that it coincides with the generating function of the generalized Bernoulli numbers (1.1).

Split the sum as $G(z) = G_1(z) + G_2(z)$, where

$$(3.2) \quad G_1(z) = \sum_{n=0}^{p-1} B_n^{(p)} \frac{z^n}{n!} \text{ and } G_2(z) = \sum_{n=p}^{\infty} B_n^{(p)} \frac{z^n}{n!}.$$

Observe that

$$\begin{aligned} G_2(z) &= \sum_{n=p}^{\infty} \frac{(-1)^{p-1}}{(p-1)!} \frac{n!}{(n-p)!} \beta^{n-p+1} (1+\beta) \cdots ((p-1)+\beta) \frac{z^n}{n!} \\ &= \frac{(-1)^{p-1}}{(p-1)!} \beta (1+\beta) \cdots (p-1+\beta) \sum_{n=p}^{\infty} \frac{n!}{(n-p)!} \beta^{n-p} \frac{z^n}{n!} \\ &= \frac{(-1)^{p-1}}{(p-1)!} (-1)^p \sum_{k=1}^p s_k^{(p)} (-1)^k z^p f_k(z) \end{aligned}$$

with

$$(3.3) \quad f_k(z) = \sum_{n=p}^{\infty} \frac{B_{n-p+k}}{(n-p)!(n-p+k)!} z^{n-p}.$$

The $(k-1)$ -st antiderivative of $f_k(z)$, denoted by $g_k(z)$, is

$$\begin{aligned} g_k(z) &= \sum_{n=p}^{\infty} \frac{B_{n-p+k}}{(n-p+k)!} z^{n-p+k-1} \\ &= z^{-1} \sum_{\ell=k}^{\infty} \frac{B_{\ell}}{\ell!} z^{\ell} \\ &= \frac{1}{z} \left[\frac{z}{e^z - 1} - \sum_{\ell=0}^{k-1} \frac{B_{\ell}}{\ell!} z^{\ell} \right], \end{aligned}$$

therefore

$$\begin{aligned} f_k(z) &= \left(\frac{d}{dz} \right)^{k-1} \frac{1}{e^z - 1} - \left(\frac{d}{dz} \right)^{k-1} \frac{1}{z} \\ &= \left(\frac{d}{dz} \right)^{k-1} \frac{1}{e^z - 1} + \frac{(-1)^k (k-1)!}{z^k}. \end{aligned}$$

This gives

$$\begin{aligned} G_2(z) &= -\frac{z^p}{(p-1)!} \sum_{k=1}^p s_k^{(p)} (-1)^k f_k(z) \\ &= -\frac{z^p}{(p-1)!} \sum_{k=1}^p s_k^{(p)} (-1)^k \left(\frac{d}{dz} \right)^{k-1} \left[\frac{1}{e^z - 1} \right] - \frac{z^p}{(p-1)!} \sum_{k=1}^p s_k^{(p)} \frac{(k-1)!}{z^k}. \end{aligned}$$

On the other hand,

$$\begin{aligned}
G_1(z) &= \sum_{n=0}^{p-1} B_n^{(p)} \frac{z^n}{n!} \\
&= \sum_{n=0}^{p-1} \frac{s_{p-n}^{(p)}}{\binom{p-1}{n}} \frac{z^n}{n!} \\
&= \frac{1}{(p-1)!} \sum_{n=0}^{p-1} s_{p-n}^{(p)} (p-1-n)! z^n \\
&= \frac{1}{(p-1)!} \sum_{k=1}^p s_k^{(p)} (k-1)! z^{p-k}.
\end{aligned}$$

This sum cancels the second term in the expression for $G_2(z)$. Hence

$$(3.4) \quad G(z) = G_1(z) + G_2(z) = -\frac{z^p}{(p-1)!} \sum_{k=1}^p s_k^{(p)} (-1)^k \left(\frac{d}{dz}\right)^{k-1} \left[\frac{1}{e^z - 1}\right].$$

Using (2.9) this gives

$$(3.5) \quad G(z) = -\frac{(-z)^p}{(p-1)!} \left((p-1) + \frac{d}{dz} \right) \cdots \left(1 + \frac{d}{dz} \right) \left[\frac{1}{e^z - 1} \right].$$

The next lemma simplifies this expression. Its proof by induction is elementary, so it is omitted.

Lemma 3.1. *For $n \geq 1$, the identity*

$$(3.6) \quad \frac{(-1)^n}{n!} \left(n + \frac{d}{dz} \right) \left(n-1 + \frac{d}{dz} \right) \cdots \left(1 + \frac{d}{dz} \right) \frac{1}{e^z - 1} = \frac{1}{(e^z - 1)^{n+1}}$$

holds.

Replacing in (3.5) produces

$$(3.7) \quad G(z) = -\frac{(-z)^p}{(p-1)!} \frac{(p-1)!}{(-1)^{p-1}} \frac{1}{(e^z - 1)^p} = \left(\frac{z}{e^z - 1} \right)^p,$$

which is the generating function of the generalized Bernoulli numbers. This proves both Lucas's formula for $B_n^{(p)}$ with $n \geq p$ and the expression (2.10) for $0 \leq p \leq n-1$.

4. LUCAS'S FORMULA VIA RECURRENCES

The numbers $B_n^{(p)}$ satisfy the recurrence

$$(4.1) \quad pB_n^{(p+1)} = (p-n)B_n^{(p)} - pnB_{n-1}^{(p)}.$$

Lucas's formula for $B_n^{(p)}$ is now established by showing that the numbers defined by (2.4) satisfy the same recurrence.

Start with

$$(p-n)B_n^{(p)} - pnB_{n-1}^{(p)} = (p-n)\frac{(-1)^{p-1}n!}{(p-1)!(n-p)!}\beta^{n-p}\prod_{k=0}^{p-1}(k+\beta) - pn\frac{(-1)^{p-1}n!}{(p-1)!(n-p-1)!}\beta^{n-1-p}\prod_{k=0}^{p-1}(k+\beta),$$

and write it as

$$\begin{aligned} (p-n)B_n^{(p)} - pnB_{n-1}^{(p)} &= \frac{(-1)^{p-1}n!}{(p-1)!(n-p-1)!}\beta^{n-1-p}\left[-\prod_{k=0}^{p-1}(k+\beta) - p\beta\prod_{k=0}^{p-1}(k+\beta)\right] \\ &= \frac{(-1)^p n!}{(p-1)!(n-p-1)!}\beta^{n-1-p}(p+\beta)\prod_{k=0}^{p-1}(k+\beta) \\ &= p\frac{(-1)^p}{p!}\frac{n!}{(n-p-1)!}\beta^{n-1-p}\prod_{k=0}^p(k+\beta) \\ &= pB_n^{(p+1)}. \end{aligned}$$

To conclude the result, it suffices to check that the initial conditions match. This is clear, since

$$(4.2) \quad B_n^{(1)} = \frac{n!}{(n-1)!}\beta^n = n\beta^n = n\frac{B_n}{n} = B_n.$$

This establishes Lucas's formula for the generalized Bernoulli numbers.

5. A NEW PROOF OF DILCHER'S FORMULA

This section analyzes the sum

$$(5.1) \quad S_N(n) := \sum \binom{2n}{2j_1, 2j_2, \dots, 2j_N} B_{2j_1} B_{2j_2} \cdots B_{2j_N},$$

using Lucas's expression for the generalized Bernoulli numbers $B_n^{(p)}$. An alternative formulation is presented.

Proposition 5.1. *The sum $S_N(n)$ is given by*

$$(5.2) \quad S_N(n) = \sum_{k=0}^N \frac{(2n)!}{(2n-k)!} 2^{-k} \binom{N}{k} B_{2n-k}^{(N-k)}$$

for $2n > N$.

Proof. The umbral method [7] shows that the sum $S_N(n)$ is given by

$$(5.3) \quad S_N(n) = \frac{1}{2^N} (\epsilon_1 B_1 + \cdots + \epsilon_N B_N)^{2n}$$

with $\epsilon_j = \pm 1$. Introduce the notation

$$(5.4) \quad Y_{2n}^{(M,N)} = (-B_1 - \cdots - B_M + B_{M+1} + \cdots + B_N)^{2n}$$

where there are M minus signs and $N - M$ plus signs. Thus,

$$(5.5) \quad S_N(n) = \frac{1}{2^N} \sum_{M=0}^N \binom{N}{M} Y_{2n}^{(M,N)}.$$

The next step uses the famous umbral identity

$$(5.6) \quad f(-B) = f(B) + f'(0)$$

(see Section 2 of [3] for details) to obtain

$$(5.7) \quad Y_{2n}^{(M,N)} = Y_{2n}^{(M-1,N)} + 2nY_{2n-1}^{(M-1,N-1)}.$$

This may be written as

$$(5.8) \quad Q_{2n}^{(M)} = Q_{2n}^{(M-1)} + 2nQ_{2n-1}^{(M-1)},$$

where $Q_j^M = Y_j^{(M,P+j)}$ and $P = N - 2n$. Then (5.8) is easily solved to produce

$$(5.9) \quad Q_{2n}^{(M)} = \sum_{k=0}^M \binom{M}{k} \frac{(2n)!}{(2n-k)!} Q_{2n-k}^{(0)}.$$

Since the initial condition is

$$(5.10) \quad Q_{2n-k}^{(0)} = Y_{2n-k}^{(0,N-k)} = B_{2n-k}^{(N-k)},$$

it follows that

$$(5.11) \quad Y_{2n}^{(M,N)} = \sum_{k=0}^M \binom{M}{k} \frac{(2n)!}{(2n-k)!} B_{2n-k}^{(N-k)}.$$

Replacing in (5.5) yields

$$\begin{aligned} S_N(n) &= \frac{1}{2^N} \sum_{M=0}^N \binom{N}{M} Y_{2n}^{(M,N)} \\ &= \frac{1}{2^N} \sum_{M=0}^N \binom{N}{M} \sum_{k=0}^M \binom{M}{k} \frac{(2n)!}{(2n-k)!} B_{2n-k}^{(N-k)} \\ &= \frac{1}{2^N} \sum_{k=0}^N \frac{(2n)!}{(2n-k)!} B_{2n-k}^{(N-k)} \sum_{M=0}^N \binom{M}{k} \binom{N}{M}. \end{aligned}$$

Now use the basic identity

$$(5.12) \quad \sum_{M=0}^N \binom{M}{k} \binom{N}{M} = \sum_{M=k}^N \binom{M}{k} \binom{N}{M} = 2^{N-k} \binom{N}{k}$$

to obtain the result. \square

Lucas's identity for generalized Bernoulli numbers is now used to obtain a second expression for the sum $S_N(n)$.

Proposition 5.2. *For $2n > N$, the sum $S_N(n)$ is given by*

$$(5.13) \quad S_N(n) = \frac{(2n)!}{(2n-N)!} \beta^{2n-N+1} \sum_{\ell=0}^{N-1} \binom{N}{\ell+1} \frac{(-1)^\ell (\beta+1)_\ell}{2^{N-1-\ell} \ell!}.$$

Proof. Using the Pochhammer symbol

$$(5.14) \quad (\beta+1)_{p-1} = \frac{\Gamma(\beta+p)}{\Gamma(\beta+1)} = (\beta+1) \cdots (\beta+p-1)$$

Lucas's formula (2.4) is stated in the form

$$(5.15) \quad B_n^{(p)} = \frac{(-1)^{p-1}}{(p-1)!} \frac{n!}{(n-p)!} \beta^{n-p+1} (\beta+1)_{p-1}.$$

Using Proposition 5.1 and $B_n^{(0)} = \delta_n$ so that $B_{2n-N}^{(0)} = 0$ since $2n > N$, it follows that

$$\begin{aligned} S_N(n) &= \sum_{k=0}^{N-1} \frac{(2n)!}{(2n-k)!} 2^{-k} \binom{N}{k} \frac{(-1)^{N-k-1}}{(N-k-1)!} \frac{(2n-k)!}{(2n-N)!} \beta^{2n-N+1} (\beta+1)_{N-k-1} \\ &= \frac{(2n)!}{(2n-N)!} \beta^{2n-N+1} \sum_{k=0}^{N-1} 2^{-k} \binom{N}{k} \frac{(-1)^{N-k-1}}{(N-k-1)!} (\beta+1)_{N-k-1} \end{aligned}$$

that reduces to the stated form. □

To obtain a hypergeometric form of the sum $S_N(n)$, observe that

$$(5.16) \quad N(1-N)_\ell = (-1)^\ell \frac{N!}{(N-\ell-1)!}$$

and $(2)_\ell = (\ell+1)!$ give

$$(5.17) \quad (-1)^\ell \binom{N}{\ell+1} = N \frac{(1-N)_\ell}{(2)_\ell},$$

and the following result follows from Proposition 5.2.

Proposition 5.3. *The hypergeometric form of the sum $S_N(n)$ is given by*

$$(5.18) \quad S_N(n) = \frac{(2n)!}{(2n-N)!} \beta^{2n-N+1} 2^{1-N} N {}_2F_1 \left(\begin{matrix} 1-N, & 1+\beta \\ & 2 \end{matrix} \middle| 2 \right).$$

The final form of the sum $S_N(n)$ involves the Meixner-Pollaczek polynomials defined by

$$(5.19) \quad P_n^{(\lambda)}(x; \phi) = \frac{(2\lambda)_n}{n!} e^{n\phi} {}_2F_1 \left(\begin{matrix} -n, & \lambda + ix \\ & 2\lambda \end{matrix} \middle| 1 - e^{-2i\phi} \right).$$

Choosing $\lambda = 1$ and $\phi = \pi/2$ gives the next result.

Theorem 5.1. *The sum $S_N(n)$ is given by*

$$(5.20) \quad S_N(n) = \frac{(2n)!}{(2n-N)!} \frac{1}{(2i)^{N-1}} \beta^{2n-N+1} P_{N-1}^{(1)} \left(-i\beta; \frac{\pi}{2} \right).$$

Some examples are presented next.

Example 5.4. *The Meixner-Pollaczek polynomial*

$$(5.21) \quad P_2^{(1)}\left(x; \frac{\pi}{2}\right) = 2x^2 - 1$$

gives

$$\begin{aligned} S_3(n) &= \frac{(2n)!}{(2n-3)!} \times (-1/4)\beta^{2n-2}(-2\beta^2 - 1) \\ &= \frac{(2n)(2n-1)(2n-2)}{4} \left[2\frac{B_{2n}}{2n} + \frac{B_{2n-2}}{2n-2} \right] \\ &= (2n-1)(n-1)B_{2n} + \frac{1}{2}n(2n-1)B_{2n-2}, \end{aligned}$$

which coincides with [2, eq. (2.6)].

Example 5.5. *The Meixner-Pollaczek of degree 3 is*

$$(5.22) \quad P_3^{(1)}\left(x; \frac{\pi}{2}\right) = \frac{4}{3}(-2x + x^3)$$

that produces

$$\begin{aligned} S_3(n) &= \frac{(2n)!}{(2n-4)!} \frac{1}{(2i)^3} \beta^{2n-3} \frac{4}{3} (2i\beta + i\beta^3) \\ &= -\frac{1}{3}(2n-1)(n-1)(2n-3)B_{2n} - \frac{1}{3}(2n)(2n-1)(2n-3)B_{2n-2}, \end{aligned}$$

which coincides with [2, eq. (2.7)].

The next step is to establish a correspondence between the *Dilcher coefficients* $b_k^{(N)}$ in (1.6) and the coefficients $p_k^{(n)}$ in

$$(5.23) \quad P_n^{(1)}(x; \pi/2) = \sum_{k=0}^n p_k^{(n)} x^k$$

the Meixner-Pollaczek polynomials. In particular, it is shown that the recurrence (1.7) is a consequence of the classical three terms recurrence for orthogonal polynomials.

Theorem 5.2. *The coefficients $b_k^{(N)}$ defined in (1.6) and the coefficients $p_k^{(n)}$ are related by*

$$(5.24) \quad b_k^{(N)} = \frac{(-1)^{N-1-k}}{2^{N-1}} p_{N-1-2k}^{(N-1)}.$$

The recurrence relation (1.7) is equivalent to the three-terms recurrence

$$(5.25) \quad (n+1)P_{n+1}^{(1)}\left(x; \frac{\pi}{2}\right) - 2xP_n^{(1)}\left(x; \frac{\pi}{2}\right) + (n+1)P_{n-1}^{(1)}\left(x; \frac{\pi}{2}\right) = 0.$$

satisfied by the Meixner-Pollaczek polynomials.

Proof. The Meixner-Pollaczek polynomials are orthogonal, hence they satisfy a three-terms recurrence. The specific form for this family in (5.25) appears in [6, Chapter 18]. In terms of its coefficients $p_k^{(n)}$ this is expressed as

$$(5.26) \quad (n+1)p_k^{(n+1)} - 2p_{k-1}^{(n)} + (n+1)p_k^{(n-1)} = 0.$$

Comparing the two expressions for $S_N(n)$ in (1.6) and (5.20) gives (5.24). This is equivalent to

$$(5.27) \quad p_\ell^{(N-1)} = 2^{N-1} t^{N-1+\ell} b_{\frac{1}{2}(N-1-\ell)}^{(N)}.$$

Replacing in (5.26) and simplifying yields (1.7). □

Theorem 2 in [2], stated below, may be proven along the same lines of the proof of Theorem 2.2. Details are omitted.

Theorem 5.3. *If $2n \leq N - 1$, then*

$$(5.28) \quad \begin{aligned} S_N(n) &= (-1)^n \frac{(2n)!(N-2n-1)!}{2^{N-1}} p_{N-2n-1}^{(N-1)} \\ &= (-1)^{N-1} (2n)!(N-2n-1)! b_n^{(N)}. \end{aligned}$$

Acknowledgments. The work of the first author was partially funded by NSF-DMS 1112656.

REFERENCES

- [1] L. Carlitz. Note on Nörlund polynomial $B_n^{(z)}$. *Proc. Amer. Math. Soc.*, 11:452–455, 1960.
- [2] K. Dilcher. Sums of products of Bernoulli numbers. *Journal of Number Theory*, 60:23–41, 1996.
- [3] A. Dixit, V. Moll, and C. Vignat. The Zagier modification of Bernoulli numbers and a polynomial extension. Part I. *The Ramanujan Journal*, To appear, 2014.
- [4] E. Lucas. Sur les congruences des nombres Euleriens et des coefficients différentiels des fonctions trigonometriques, suivant un module premier. *Bull. Soc. Math. France*, 6:49–54, 1878.
- [5] N. E. Nörlund. *Vorlesungen über Differenzen-Rechnung*. Berlin, 1924.
- [6] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, editors. *NIST Handbook of Mathematical Functions*. Cambridge University Press, 2010.
- [7] S. Roman. *The Umbral Calculus*. Dover, New York, 1984.

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LA 70118
E-mail address: vhm@tulane.edu

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LA 70118
E-mail address: cvignat@math.tulane.edu